

TRUNCATIONS OF LEVEL 1 OF ELEMENTS IN THE LOOP GROUP OF A REDUCTIVE GROUP

EVA VIEHMANN

ABSTRACT. We generalize the notion of Ekedahl-Oort strata to elements in the loop group of any connected reductive group, and call the resulting discrete invariant the truncation of level 1 of the element. We give conditions for the Newton points occurring among the elements of a given truncation of level 1 and especially for the generic Newton point in a given truncation stratum. We prove that truncation strata are locally closed and give a description of the closure of each stratum. We also translate our results back to the original Ekedahl-Oort stratification of the reduction modulo p of Shimura varieties.

1. INTRODUCTION

Let k be an algebraically closed field of characteristic p . Let L be either $k((t))$ or $\text{Quot}(W(k))$ and let $\mathcal{O}$ be the valuation ring. We denote by $\sigma : x \mapsto x^{p^r}$ the Frobenius of k over $\mathbb{F}_{p^r}$ for some fixed r and also the associated Frobenius of L over $F = \mathbb{F}_{p^r}((t))$ resp. $\mathbb{Q}_{p^r}$. Let $\mathcal{O}_F$ be the valuation ring of F . We denote the uniformizer t or p of $\mathcal{O}_F$ by ϵ .

Let G be a connected reductive group over $\mathcal{O}_F$. Then G is unramified, i. e. quasi-split and split over an unramified extension of $\mathcal{O}_F$. Let B be a Borel subgroup of G and let T be a maximal torus contained in B . Let $K = G(\mathcal{O})$ and let I be the inverse image of $B(k)$ under the projection $K \rightarrow G(k)$. Let K_1 be the kernel of the projection $K \rightarrow G(k)$. Let $W = N_T(L)/T(L)$ denote the (absolute) Weyl group of T in G where N_T denotes the normalizer of T . Let $\widetilde{W} = N_T(L)/T(\mathcal{O}) \cong W \ltimes X_*(T)$ denote the affine Weyl group. For each $w \in W$ we choose a representative in $N_T(\mathcal{O})$. We denote this representative by the same letter as the element itself. If M is a Levi subgroup of G containing T let W_M be the Weyl group of M and denote by ${}^M W$ resp. ${}^M \widetilde{W}$ the set of elements x of W resp. $\widetilde{W}$ that are shortest representatives of their coset $W_M x$. Similarly, W^M denotes the set of elements x that are the shortest representatives of their cosets $x W_M$ and accordingly for $\widetilde{W}$.

For $b \in G(L)$ we call $\{g^{-1}b\sigma(g) \mid g \in K\}$ the K - σ -conjugacy class of b , and $[b] = \{g^{-1}b\sigma(g) \mid g \in G(L)\}$ the σ -conjugacy class of b .

The aim of this article is to classify and describe the K - σ -conjugacy classes of elements of $K_1 \backslash G(L) / K_1$.

1.1. Comparison with Ekedahl-Oort strata. Let X be a p -divisible group over an algebraically closed field k of characteristic p . Then the Dieudonné module of X is a pair $(\mathbf{M}, F)$ where $\mathbf{M}$ is a free $W(k)$ -module of rank equal to the height h of X and where $F : \mathbf{M} \rightarrow \mathbf{M}$ is a σ -linear homomorphism. Here $W(k)$ is the ring of Witt vectors of k and σ denotes the Frobenius of $W(k)$ over $\mathbb{Z}_p$. Choosing a basis for $\mathbf{M}$ we can write $F = b\sigma$ for some $b \in GL_h(W(k)[1/p])$. A change of the basis amounts to σ -conjugating b by an element of $GL_h(W(k)) = K$. As b is induced by the Dieudonné module $(\mathbf{M}, F)$, we have $b \in Kp^\mu K$ for some minuscule $\mu \in X_*(T)$. Similarly, polarized p -divisible groups or abelian varieties lead to elements $b \in GSp_{2n}(W(k)[1/p])$ for n equal to the dimension of the p -divisible group.

2000 *Mathematics Subject Classification.* 14G35, 20G25, 14K10.

Key words and phrases. Ekedahl-Oort stratification, σ -conjugacy class, loop group.

In [O1] Oort shows that one obtains a discrete invariant of X (the so-called Ekedahl-Oort invariant) by considering the p -torsion points $X[p]$, or equivalently by studying the reduction modulo p of the Dieudonné module $\mathbf{M}$ together with the two maps induced by $F : \mathbf{M} \rightarrow \mathbf{M}$ and $V = pF^{-1} : \mathbf{M} \rightarrow \mathbf{M}$. In terms of the element b , this corresponds to considering the K_1 -double coset. In other words, this situation is analogous to the above in the special case $G = GL_h$ or GSp_{2n} and μ minuscule for $\mathcal{O} = W(k)$. A classification of the Ekedahl-Oort invariant (and more generally of $G(k)$ -orbits on a certain variety associated to a reductive group G over $\mathbb{Z}_p$ together with a fixed Levi subgroup that is also defined over $\mathbb{Z}_p$) using a similar subset of the Weyl group of the associated reductive group as we use has been given by Moonen and Wedhorn in [MW].

The classification of the K - σ -conjugacy classes of elements of $K_1 \backslash G(L) / K_1$ is given by the following theorem (which we prove in Section 2). The second part of the theorem establishes a relation between the subdivisions of $K_1 \backslash G(L) / K_1$ according to K - σ -conjugacy classes and according to Iwahori-double cosets. Let us first introduce some notation. For a dominant $\mu \in X_*(T)$ let M_μ be the centralizer of μ and let ${}^\mu W = \sigma^{-1}({}^{M_\mu} W)$. Let $x_\mu = w_0 w_{0,\mu}$ where w_0 denotes the longest element of W and where $w_{0,\mu}$ is the longest element of W_{M_μ} . Let $\tau_\mu = x_\mu \epsilon^\mu$ where ϵ^μ is the image of ϵ under $\mu : \mathbb{G}_m \rightarrow T$. Then τ_μ is the shortest element of $W \epsilon^\mu W$.

Theorem 1. *Let $b \in G(L)$. Let $\mu \in X_*(T)$ be the unique dominant element with $b \in K \epsilon^\mu K$.*

- (1) *There is a unique $w \in {}^\mu W$ such that the K - σ -conjugacy class of b contains an element of $K_1 w \tau_\mu K_1$.*
- (2) *Let $w \in {}^\mu W$. Then each element of $I w \tau_\mu I$ is I - σ -conjugate to an element of $K_1 w \tau_\mu K_1$.*

Definition 2. Let $b \in G(L)$. The pair (w, μ) as in Theorem 1 is called the *truncation of level 1* of b .

Definition 3. Let $\mu \in X_*(T)$ be dominant, let $w \in {}^\mu W$ and assume that $\text{char}(F) = p$. Then let $S_{w,\mu}$ be the reduced subscheme of the loop group of $G_{\mathbb{F}_{p^r}}$ such that $S_{w,\mu}(k)$ consists of those $g \in G(k((t)))$ whose truncation of level 1 is (w, μ) .

Here the loop group of a group G over $\mathbb{F}_{p^r}$ is the group ind-scheme representing the functor on $\mathbb{F}_{p^r}$ -algebras $R \mapsto G(R((t)))$.

For $F = \mathbb{Q}_{p^r}$ it is not clear how to define an ind-scheme having $G(L)$ as its set of k -valued points. However one can study the stratifications induced on the reduction modulo p of certain Shimura varieties (compare 1.1). The main part of this paper is concerned with elements of $G(L)$ for both cases or, whenever a scheme structure is involved, the equal characteristic case. In Section 6 we discuss applications to Shimura varieties.

We show in Section 5 that the $S_{w,\mu}$ are bounded locally closed subschemes of the loop group and that the closure of each stratum is a union of finitely many strata. The following criterion generalizes a corresponding result for Ekedahl-Oort strata by Wedhorn [W] to our situation.

Theorem 4. *$S_{w',\mu'} \subseteq \overline{S_{w,\mu}}$ if and only if there is a $\tilde{w} \in W$ with $\tilde{w} w' \tau_{\mu'} \sigma(\tilde{w})^{-1} \leq w \tau_\mu$ with respect to the Bruhat order.*

1.2. Truncations of level 1 and σ -conjugacy classes. One interesting open question about Ekedahl-Oort strata is to determine which Newton polygons occur in a given Ekedahl-Oort stratum. Recently progress towards answering this question has been made in two ways. In a series of papers [Ha1], [Ha2], [Ha3] Harashita proves a conjecture of Oort ([O2], 6.9) giving a characterization of the generic Newton polygon in each Ekedahl-Oort stratum in the moduli space of principally polarized abelian varieties. Besides, Görtz, Haines, Kottwitz and Reuman [GHR] study the intersections between Iwahori double cosets in the loop group of a reductive group and σ -conjugacy classes. In this paper we generalize the necessary results from [GHR] from split to unramified reductive groups. Together with Theorem 1(2) this allows to deduce

similar conditions for the intersections between the truncation strata and σ -conjugacy classes. Especially, we obtain a generalization of Harashita's theorem to the loop group of any connected reductive group.

To formulate this generalization let us review Kottwitz's classification [K1] of the set $B(G)$ of σ -conjugacy classes of elements $b \in G(L)$ which generalizes the notion of Newton polygons (compare also [RR], Section 1 for a more complete review of these results). Each σ -conjugacy class is determined by two invariants. One of them is given by a map $\kappa_G : B(G) \rightarrow \pi_1(G)_\Gamma$ where $\pi_1(G)$ is the quotient of $X_*(T)$ by the coroot lattice and where Γ is the absolute Galois group of F . There is the following explicit description for κ_G . Let $b \in G(L)$ and let $r_B(b) = \mu \in X_*(T)$ be the unique element such that $b \in Ke^\mu U(L)$ where U is the unipotent radical of B . Then $\kappa_G(b)$ is the image of μ under the canonical projection from $X_*(T)$ to $\pi_1(G)_\Gamma$. The second invariant is the so-called the Newton point $\nu = \nu_b$ of b , an element of $(X_*(T)_\mathbb{Q}/W)^\Gamma$, the set of Γ -invariant W -orbits on $X_*(T) \otimes \mathbb{Q}$. This invariant is the direct analog of the usual Newton polygon classifying F -isocrystals over an algebraically closed field.

In [RR] Rapoport and Richartz define a partial order on $B(G)$ given by $[b] \preceq [b']$ if and only if $\kappa_G(b) = \kappa_G(b')$ and $\nu_b \preceq \nu_{b'}$. Here the second condition means that $\tilde{\nu}_{b'} - \tilde{\nu}_b$ is a linear combination of positive roots with coefficients in $\mathbb{Q}_{\geq 0}$ where $\tilde{\nu}_{b'}$ and $\tilde{\nu}_b$ are dominant representatives of the two orbits (compare Lemma 2.2 of loc. cit.). Their Theorem 3.6 shows that for each $[b]$, the union of all σ -conjugacy classes which are less or equal to $[b]$ is closed in the loop group. More precisely, they show a corresponding statement over a field F of mixed characteristic. The function field analog can be shown in a similar, but slightly easier way using properties of the affine Grassmannian, compare [HV], Theorem 6.3.

In Section 3 we construct for each σ -conjugacy class $[b]$ a unique so-called minimal truncation type (w_b, μ_b) . It satisfies $w_b \tau_{\mu_b} \in [b]$ and a technical property (see Definition 7) which ensures that $Iw_b \tau_{\mu_b} I$ is even K - σ -conjugate to $w_b \tau_{\mu_b}$. The element $w_b \tau_{\mu_b}$ is also called the minimal element in the given class. This is a generalization of the notion of minimal p -divisible groups (as in [O3]) to our context (compare Section 6). The following theorem is the function field analog for general G and μ of Oort's conjecture and also justifies the name minimal truncation type.

Theorem 5. *Let $b \in G(L)$, and let (w, μ) be its truncation of level 1. Let $w_b \tau_{\mu_b}$ be the minimal element in the σ -conjugacy class of b . Then $w_b \tau_{\mu_b} \in \overline{S_{w, \mu}}$.*

From Theorem 5, one can easily deduce the following corollary which gives a characterization of the generic Newton point in a given stratum $S_{w, \mu}$.

Corollary 6. *Let $[b]$ be the generic σ -conjugacy class in $S_{w, \mu}$ for some $w \in {}^\mu W$. Then $[b]$ is the maximal element in the set of σ -conjugacy classes of minimal elements $w' \tau_{\mu'} \in \widetilde{W}$ such that $S_{w', \mu'} \subseteq \overline{S_{w, \mu}}$. This is also the same as the maximal class $[x]$ among all $x \in \widetilde{W}$ with $x \leq w \tau_\mu$.*

Here we use that $S_{w, \mu}$ is irreducible, so it contains a unique generic σ -conjugacy class.

In the last section we give a translation of our results and their proofs back to the case of mixed characteristic. In particular, one obtains Oort's conjecture (as shown by Harashita) as well as generalizations to other cases of good reduction of Shimura varieties of PEL type.

While finishing this article Harashita published a preprint [Ha4] in which he generalizes his proof of Oort's conjecture to also obtain an analog for some catalog of p -divisible groups in the non-polarized case.

Acknowledgement. I am grateful to Ulrich Görtz, Xuhua He, Torsten Wedhorn, and Chia-Fu Yu for helpful discussions. Part of this paper was written during a stay at the Academia Sinica in Taipeh which was supported by the Academia Sinica and by the Hausdorff Research Center, Bonn. I thank the Academia Sinica for its hospitality, support and excellent working conditions.

2. TRUNCATIONS OF LEVEL 1

The goal of this section is to prove Theorem 1, in particular we allow both the function field case and the case of mixed characteristic. The proof follows a strategy by Bédard, [B].

Proof of Theorem 1. Let $b \in G(L)$. Then there is a unique dominant $\mu \in X_*(T)$ with $b \in K\epsilon^\mu K$ and b is K - σ -conjugate to an element of the form $b_0 x_\mu \epsilon^\mu = b_0 \tau_\mu$ with $b_0 \in K$.

To show (1) we use induction on i to show that there exist a sequence of elements $u_i \in W$ and two sequences of Levi subgroups M_i, M'_i of G with the following properties:

- a) $M_0 = M'_0 = G$,
 $M'_1 = x_\mu M_\mu x_\mu^{-1}$ and $M_1 = \sigma^{-1}(M_\mu)$,
 $M_i = M'_{i-1} \cap \sigma^{-1}(u_{i-1}^{-1} M_\mu u_{i-1})$, and
 $M'_i = M'_1 \cap x_\mu \sigma(u_{i-1} M'_{i-1} u_{i-1}^{-1}) x_\mu^{-1} = x_\mu \sigma(u_{i-1} M_i u_{i-1}^{-1}) x_\mu^{-1}$ for $i > 0$.
- b) $u_0 = 1$ and
 $u_i = u_{i-1} \delta_i$ for $i > 0$ for some $\delta_i \in W_{M'_{i-1}}$ which is the shortest representative of $W_{M_i} \delta_i W_{M'_i}$.
- c) b is K - σ -conjugate to an element of $K_1 u_i b_i \tau_\mu K_1$ with $b_i \in M'_i(\mathcal{O})$.
- d) $u_i W_{M'_i} \subseteq W$ as in c) is uniquely determined by the K - σ -conjugacy class of b in $K_1 \backslash G(L) / K_1$.
- e) Let $u_i b_i \tau_\mu$ be as in c). Then $u_i b'_i \tau_\mu$ with $b'_i \in M'_i(\mathcal{O})$ is in the K - σ -conjugacy class of b in $K_1 \backslash G(L) / K_1$ if and only if modulo K_1 the two elements b_i, b'_i are in the same $M'_{i+1}(k)$ - σ -conjugacy class in $(U_{P_{i+1}}(k) \cap M'_i(k)) \backslash M'_i(k) / (U_{\overline{P}'_{i+1}}(k) \cap M'_i(k))$. Here P_{i+1} is the parabolic with Levi subgroup M_{i+1} which contains B , and $P'_{i+1} = x_\mu \sigma(u_i P_{i+1} u_i^{-1}) x_\mu^{-1}$. Furthermore U_P denotes the unipotent radical of a parabolic P .

One easily checks that a) and b) imply that $M_i \supseteq M_{i+1}$, and $M'_i \supseteq M'_{i+1}$ for all i .

For $i = 0$, d) is obvious and c) has been shown above. For e) note that modulo K_1 , the effect of σ -conjugation of $b_0 \tau_\mu$ by $g \in U_{P_1}(\mathcal{O})$ is the same as left multiplication of b_0 by g . Right multiplication of b_0 by an element of $U_{\overline{P}'_1}(\mathcal{O})$ corresponds to multiplying $b_0 \tau_\mu$ by an element of $\epsilon^{-\mu} U_{\overline{P}'_1}(\mathcal{O}) \epsilon^\mu \subset K_1$. Finally if an element $g \in K$ conjugates $b_0 \tau_\mu$ into $M'_0(\mathcal{O}) \tau_\mu = K \epsilon^\mu$, then it is in $K \cap \epsilon^{-\mu} K \epsilon^\mu$. Decomposing g into factors in $U_{P_1}(\mathcal{O})$, $\epsilon^{-\mu} U_{\overline{P}'_1}(\mathcal{O}) \epsilon^\mu$ and $M_1(\mathcal{O})$ shows e).

We have to show that a)–e) for some i imply the same properties for $i + 1$. Let b_i be as in c). We decompose b_i using the Bruhat decomposition to obtain that $b_i \in P_{i+1}(\mathcal{O}) \delta_{i+1} \overline{P}'_{i+1}(\mathcal{O})$ for some δ_{i+1} as in b) and with P_{i+1} and P'_{i+1} as in e). By e) we may assume that the factors in $P_{i+1}(\mathcal{O})$ and $\overline{P}'_{i+1}(\mathcal{O})$ lie in $M_{i+1}(\mathcal{O})$ and $M'_{i+1}(\mathcal{O})$, respectively. After σ -conjugating $u_i b_i \tau_\mu$ with the factor in $u_i M_{i+1}(\mathcal{O}) u_i^{-1}$ we obtain c) for $i + 1$. Property d) follows from the uniqueness of the Bruhat decomposition together with d) and e) for i . It remains to show e). Multiplication of b_{i+1} on the left with an element of $u_{i+1}^{-1} \sigma^{-1}(U_{P_1}(\mathcal{O})) u_{i+1} \cap M'_{i+1}(\mathcal{O}) \supseteq U_{P_{i+2}}(\mathcal{O}) \cap M'_{i+1}(\mathcal{O})$ corresponds to multiplication of $u_{i+1} b_{i+1} \tau_\mu$ on the left with an element of $\sigma^{-1}(U_{P_1}(\mathcal{O})) \cap u_{i+1} M'_{i+1}(\mathcal{O}) u_{i+1}^{-1}$. Thus it does not change the K - σ -conjugacy class. Now we want to show that multiplication of b_{i+1} on the right by an element of $M'_{i+1}(\mathcal{O}) \cap x_\mu \sigma(u_{i+1} U_{\overline{P}'_{i+1}}(\mathcal{O}) u_{i+1}^{-1}) x_\mu^{-1} = M'_{i+1}(\mathcal{O}) \cap U_{\overline{P}'_{i+2}}(\mathcal{O})$ also does not change the K - σ -conjugacy class of $u_{i+1} b_{i+1} \tau_\mu$. As $M'_{i+1} \subseteq x_\mu M_\mu x_\mu^{-1}$ the multiplication of b_{i+1} corresponds to multiplying $u_{i+1} b_{i+1} \tau_\mu = u_{i+1} b_{i+1} x_\mu \epsilon^\mu$ on the right with an element of $x_\mu^{-1} M'_{i+1}(\mathcal{O}) x_\mu \cap \sigma(u_{i+1} U_{\overline{P}'_{i+1}}(\mathcal{O}) u_{i+1}^{-1})$. After σ -conjugating with this element (which is in K), we obtain an element of $u_{i+1} (U_{\overline{P}'_{i+1}}(\mathcal{O}) \cap M_{i+1}(\mathcal{O})) b_{i+1} x_\mu \epsilon^\mu$. As $b_{i+1} \in M'_{i+1}(\mathcal{O})$, the element lies in $u_{i+1} b_{i+1} (U_{\overline{P}'_{i+1}}(\mathcal{O}) \cap M_{i+1} M'_{i+1}(\mathcal{O})) x_\mu \epsilon^\mu$. Using induction and $M_{i+1} M'_{i+1} \subseteq M'_i$ we obtain that this is contained in the same class as $u_{i+1} b_{i+1} x_\mu \epsilon^\mu$. The other direction of e) is shown in the same way as for $i = 0$.

As the sequences M_i, M'_i are decreasing families of Levi subgroups, they become constant after finitely many steps. Thus for n sufficiently large, $x_\mu \sigma(u_n M'_n u_n^{-1}) x_\mu^{-1} = M'_n = M'_{n+1}$. Especially, $M_n = M'_n$. Besides, $M_n \subseteq M_1 = \sigma^{-1}(M_\mu)$. Using a variant of Lang's theorem we see that each element of $u_n M'_n(\mathcal{O}) \tau_\mu$ is $M_n(\mathcal{O})$ - σ -conjugate to $u_n \tau_\mu$. Results of Bédard [B], or Lusztig ([L], (d) in the proof of Proposition 2.4) show that b) implies that $u_n \in W$ is the shortest representative of $W_{M_1} u_n W_{M'_n}$ and thus u_n is the desired element of ${}^\mu W$.

We now prove (2). Each element of $I w \tau_\mu I$ is obviously I - σ -conjugate to some element $g \in w \tau_\mu I$. We have the Iwahori decomposition $I = N_\mu(\mathcal{O}) I_{M_\mu} K_1$ where $I_{M_\mu} = I \cap M_\mu(\mathcal{O})$ and where N_μ is the unipotent radical of $P_\mu = M_\mu B$. Applying this to the corresponding factor of g and using that $\epsilon^\mu N_\mu(\mathcal{O}) \subseteq (N_\mu(\mathcal{O}) \cap K_1) \epsilon^\mu$ we see that we may assume that $g \in w \tau_\mu I_{M_\mu}$. Thus it is I_{M_1} - σ -conjugate to an element of $I_{M_1} w \tau_\mu = I_{M_1} w x_\mu \epsilon^\mu$. As $w \in {}^\mu W = {}^{M_1} W$, conjugation by w maps positive roots in M_1 to positive roots (not necessarily in M_1). Hence we have $w^{-1} I_{M_1} w \subset I \cap (w^{-1} M_1(\mathcal{O}) w)$. Conjugation by w_0 maps all positive roots to negative roots, conjugation by $w_{0,\mu}$ maps positive roots in M_μ to negative roots in M_μ (and vice versa) and leaves positive roots in N_μ positive. Hence

$$\begin{aligned} I_{M_1} w x_\mu \epsilon^\mu &= w x_\mu ((w x_\mu)^{-1} I_{M_1} w x_\mu) \epsilon^\mu \\ &\subseteq w x_\mu (I_{M_\mu} \cap (w x_\mu)^{-1} I_{M_1} w x_\mu) \overline{N}_\mu(\mathcal{O}) \epsilon^\mu \\ &\subseteq w x_\mu (I_{M_\mu} \cap (w x_\mu)^{-1} I_{M_1} w x_\mu) \epsilon^\mu K_1. \end{aligned}$$

Iterating this argument we see that the element g is I - σ -conjugate to an element of $K_1 w x_\mu I_\infty \epsilon^\mu K_1$ where $I_\infty = I \cap \bigcap_{i \geq 0} (\text{Ad}_{(w x_\mu)^{-1} \sigma^{-1}})^i (M_\mu(\mathcal{O}))$ and where $\text{Ad}_{(w x_\mu)^{-1}}$ denotes conjugation with the given element. As I_∞ commutes with ϵ^μ and satisfies $(w x_\mu)^{-1} \sigma^{-1} (I_\infty) (w x_\mu) = I_\infty$ we can again apply a variant of Lang's theorem to obtain that each element of $K_1 w x_\mu I_\infty \epsilon^\mu K_1$ is I_∞ - σ -conjugate to an element in $K_1 w \tau_\mu K_1$. $\square$

3. FUNDAMENTAL ALCOVES

In this section we still consider both possible cases for F . We recall the definition of P -fundamental alcoves from [GHKR], 13.

Definition 7. Let P be a semistandard parabolic subgroup of G , i.e. a parabolic subgroup containing T but not necessarily B , and defined over $\mathcal{O}_F$. Let N be its unipotent radical and M the Levi factor containing T . Let $\overline{N}$ be the unipotent radical of the opposite parabolic. Let $I = I_N I_M I_{\overline{N}}$ be the corresponding Iwahori decomposition. Then an element $x \in \widetilde{W}$ defined over F is called P -fundamental if $x \in \widetilde{W}_M$, $x I_M x^{-1} = I_M$, $x I_N x^{-1} \subseteq I_N$, and $x^{-1} I_{\overline{N}} x \subseteq I_{\overline{N}}$.

Remark 8. For $x \in \widetilde{W}$ that are P -fundamental for some P , Görtz, Haines, Kottwitz, and Reuman show in [GHKR], Proposition 6.3.1 that every element of $I x I$ is I - σ -conjugate to x .

In [GHKR], Corollary 13.2.4 they show that each σ -conjugacy class $[b]$ of elements of $G(L)$ contains an element of $\widetilde{W}$ which is P -fundamental for some semistandard P .

Although their assertions are only formulated in the function field case, and for split groups, the argument carries over to our more general situation.

Lemma 9. (1) Let $x, x' \in \widetilde{W}$ be P -fundamental resp. P' -fundamental with $[x] = [x']$. Then x, x' are σ -conjugate under W , and in particular K - σ -conjugate to each other.
(2) Let $b \in G(L)$. The truncation type (w_b, μ_b) of the elements of $[b]$ which are P -fundamental for some P is as follows. Let M_ν be the centralizer of the G -dominant Newton point ν of $[b]$, a standard Levi subgroup of G defined over $\mathcal{O}_F$. The element b is σ -conjugate to an element b' of M_ν such that the M_ν -dominant Newton point of b' is equal to ν . Then $\kappa_{M_\nu}(b')$ only depends on the $G(L)$ - σ -conjugacy class $[b]$. Let

- $b_0 \in \widetilde{W}_{M_\nu}$ be the unique element with $\kappa_{M_\nu}(b_0) = \kappa_{M_\nu}(b')$ and $b_0^{-1}I_{M_\nu}b_0 = I_{M_\nu}$. Here $I_{M_\nu} = I \cap M_\nu(L)$. Then $b_0 \in [b]$ and the truncation of level 1 of b_0 is equal to (w_b, μ_b) .
- (3) Let $x \in [b] \cap \widetilde{W}$ be P -fundamental for some P . Then x is Q -fundamental for a semistandard parabolic $Q = M_Q N_Q$ if and only if $x \in \widetilde{W}_{M_Q}$, the M_Q -dominant Newton point ν_Q of x is central in M_Q , and $\langle \nu_Q, \alpha \rangle \geq 0$ for each root α of T in N_Q .

The element b_0 is also called the standard representative of $[b]$. Note that ν (and hence M_ν) are invariant under σ .

Proof. We first show (3). Let x be P -fundamental and let $P = MN$ where M is the Levi factor of P containing T . Let M' be the centralizer of the M -dominant Newton point of x . As $xI_Mx^{-1} = I_M$, the Newton point of x is central in M and hence $M \subseteq M'$. Let P' be the parabolic subgroup generated by M' and N . Let N' be its unipotent radical. Hence $P' \supseteq P$ and $N' \subseteq N$. Thus in order to show that x is P' -fundamental it is enough to verify $xI_{M'}x^{-1} = I_{M'}$. We consider the decomposition $I_{M'} = I_M I_{N \cap M'} I_{\overline{N} \cap M'}$. Each of the subgroups $M, N \cap M', \overline{N} \cap M'$ is stable under x , so we can consider each factor of $I_{M'}$ separately. For I_M the assertion is just the assumption. For $I_{N \cap M'}$ we have $x^i I_{N \cap M'} x^{-i} \subseteq I_{N \cap M'}$ for every $i > 0$. For i with $x^i \in X_*(T)$ we have equality because then $\langle \alpha, x^i \rangle = \langle \alpha, i\nu \rangle = 0$ for every root of T in M' . Considering the whole chain of containments we obtain equality for every i . A similar argument applies to $\overline{N} \cap M'$. It remains to show that if $Q \subseteq P'$ is as in the assertion, then x is Q -fundamental, but this is obvious.

Let P, P', x , and x' be as in (1). Let $P = MN$. By (3) we may assume that M is equal to the centralizer of the M -dominant Newton point of x and analogously for P' . As $[x] = [x']$, their G -dominant Newton points coincide and $\kappa_G(x) = \kappa_G(x')$. Note that the Newton point of x together with $\kappa_G(x)$ determines $\kappa_M(x)$. Hence there is a $w \in W$, invariant under σ , that conjugates I_M to $I_{M'}$, and also the values of $\kappa_M, \kappa_{M'}$ to each other. As x is uniquely determined by $\kappa_M(x)$ and the condition that $xI_Mx^{-1} = I_M$, we obtain (1).

The description of x in terms of M given above also implies (2). $\square$

Definition 10. The element $w_b \tau_{\mu_b}$ as in Lemma 9 (2) is called the minimal element of $[b]$, and (w, μ) is called the minimal truncation type of $[b]$.

4. A NON-EMPTYNESS CRITERION FOR THE INTERSECTION WITH A GIVEN σ -CONJUGACY CLASS

Remark 11. By Theorem 1 (2), $S_{w, \mu}$ has nonempty intersection with some σ -conjugacy class $[b]$ if and only if $[b] \cap Iw\tau_\mu I \neq \emptyset$. Besides, the generic σ -conjugacy class in $S_{w, \mu}$ coincides with the generic class in $Iw\tau_\mu I$.

Proposition 12. Let $b \in G(L)$ and let $x_b \in \widetilde{W} \cap [b]$ be a P -fundamental alcove for some semistandard parabolic subgroup P of G . Let M be the Levi component of P containing T . Let $x \in \widetilde{W}$ with $b \in IxI$. Then there is a $w \in {}^M W$ with $w^{-1}x_b\sigma(w) \leq x$ in the Bruhat order.

Proof. Let N be the unipotent radical of P . Let I' be an Iwahori subgroup of G of the form $I' = vIv^{-1}$ for some $v \in W$ and such that its reduction modulo t lies in $P(k)$. Let $g \in G(L)$ with $g^{-1}x_b\sigma(g) = b \in IxI$. Using the Iwasawa decomposition and the Bruhat decomposition we write $g = nmi_1wi_2$ with $n \in N(L)$, $m \in M(L)$, $i_1 \in I'$, $i_2 \in I$, and $w \in W$. By the Iwahori decomposition, $i_1 \in P(\mathcal{O})K_1$. As $w^{-1}K_1w \subseteq I$, we may assume that $i_1 = \text{id}$. Furthermore we may assume that $i_2 = \text{id}$ without changing the property $g^{-1}x_b\sigma(g) \in IxI$. Thus $g = nmw$ with $g^{-1}x_b\sigma(g) \in IxI$. Finally we may assume that w is of minimal length in its coset $W_M w$. We have

$$(1) \quad g^{-1}x_b\sigma(g) = w^{-1}m^{-1} [n^{-1}x_b\sigma(n)x_b^{-1}] x_b\sigma(mw).$$

We abbreviate the expression in the bracket, which is in $N(L)$, by $\tilde{n}$.

Let $\mathcal{N}$ be the loop group associated to N over k , i. e. the group ind-scheme representing the functor on k -algebras $R \mapsto N(R((t)))$. Let $\chi \in X_*(T)$ be central in M and such that $\langle \alpha, \chi \rangle > 0$ for every simple root α of T in N . Let

$$\begin{aligned} \phi : \mathbb{A}_k^1 \setminus \{0\} &\rightarrow \mathcal{N} \\ a &\mapsto \chi(a) \tilde{n} \chi(a)^{-1}. \end{aligned}$$

Let α be a root of T in N and let U_α denote the corresponding root subgroup. Conjugation by $\chi(a)$ maps $U_\alpha(y)$ to $U_\alpha(a^j y)$ where $j = \langle \alpha, \chi \rangle > 0$. Especially, ϕ has an extension to a morphism $\phi : \mathbb{A}_k^1 \rightarrow \mathcal{N}$ that maps 0 to id. As $\chi(a)$ is central in M ,

$$w^{-1} m^{-1} \phi(a) x_b \sigma(mw) = (w^{-1} \chi(a) w) w^{-1} m^{-1} \tilde{n} x_b \sigma(mw) (\sigma(w)^{-1} \chi(a)^{-1} \sigma(w))$$

for every $a \neq 0$. Using (1), we obtain that this is in IxI . Hence

$$w^{-1} m^{-1} \phi(0) x_b \sigma(mw) = w^{-1} m^{-1} x_b \sigma(mw) \in \overline{IxI}.$$

Let $I_M = I \cap M(L)$. The minimality property of w implies that for any positive root α of T in M the root β with $w^{-1} U_\alpha w = U_\beta$ is also positive (although not necessarily in M). As I and M are defined over $\mathcal{O}_F$, the same holds for $\sigma(w)$. Thus

$$w^{-1} I_M m^{-1} x_b \sigma(m) I_M \sigma(w) \subseteq I w^{-1} m^{-1} x_b \sigma(mw) I \subseteq \overline{IxI}.$$

As x_b is P -fundamental, it satisfies $x_b I_M x_b^{-1} = I_M$. Thus $I_M x_b I_M$ is the unique closed I_M -double coset of the form $I_M h I_M$ with $h \in M(L)$ and $\kappa_M(h) = \kappa_M(x_b)$. It is contained in the closure of any other such double coset. Applying this to $I_M m^{-1} x_b \sigma(m) I_M$ we obtain

$$w^{-1} I_M x_b I_M \sigma(w) \subseteq \overline{IxI}.$$

This implies $w^{-1} x \sigma(w) \in \overline{IxI}$. $\square$

If G is split, the first (and largest) part of the proof of Proposition 12 also follows from a non-emptiness criterion by Görtz, Haines, Kottwitz, and Reuman, [GHR], Corollary 12.1.2.

Corollary 13. *Let $[b_x]$ be the generic σ -conjugacy class in IxI for some $x \in \widetilde{W}$. Then $[b_x]$ is the largest (with respect to the order described in the introduction) among the classes $[y]$ where $y \in \widetilde{W}$ with $y \leq x$ in the Bruhat order. It is also equal to the largest the $[y]$ where $y \leq x$ is in addition of the form $y = w^{-1} z \sigma(w)$ for some P -fundamental alcove z for some semistandard $P = MN$ and where $w \in {}^M W$.*

Proof. This follows from Proposition 12 using the Grothendieck specialization theorem [RR], Theorem 3.6 together with the fact that the generic σ -conjugacy classes of IxI and $\overline{IxI}$ coincide. $\square$

Proof of Theorem 5 and Corollary 6. Theorem 5 and Corollary 6 follow from Proposition 12 and Corollary 13 by Remark 11. $\square$

5. CLOSURE RELATIONS

Lemma 14. (1) *Each $S_{w,\mu}$ is bounded and admissible, smooth and irreducible.*
(2) *The closure $\overline{S_{w,\mu}}$ of $S_{w,\mu}$ is a union of finitely many strata $S_{w',\mu'}$.*
(3) *$S_{w,\mu}$ is locally closed.*
(4) *$g_0 \in G(L)$ is in $\overline{S_{w,\mu}}$ if and only if it is K - σ -conjugate to an element of $\overline{Iw\tau_\mu I}$.*

Here by admissible we mean that the subscheme is invariant under some bounded open subgroup of the loop group. We call an admissible subscheme X smooth resp. irreducible if the same holds for $X/\bar{K}$ for some (and then every) such subgroup.

Proof. The stratum is bounded because it is contained in $Kt^\mu K$, and admissible because it is invariant under K_1 . The other two assertions of (1) follow as $S_{w,\mu}/K_1$ is a single K - σ -orbit. For the second assertion note that $\overline{S_{w,\mu}}$ is invariant under K - σ -conjugation and under multiplication by K_1 on both sides. Thus it is a union of strata. The union is finite because each of these strata $S_{w',\mu'}$ has to satisfy $\mu' \preceq \mu$. (3) follows from (1) and (2). In (4) the second condition obviously implies the first. Now let $g_0 \in \overline{S_{w,\mu}}$. Thus there is a $g \in G(k[[z]]((t)))$ such that its reduction modulo z is equal to g_0 and such that its image g_η in $G(k((z))((t)))$ is in $S_{w,\mu}$. Hence there is an $h \in G(k((z))^{\text{alg}}[[t]])$ with $h^{-1}g_\eta\sigma(h) \in K_{1,k((z))^{\text{alg}}}w\tau_\mu K_{1,k((z))^{\text{alg}}}$. Here $k((z))^{\text{alg}}$ denotes an algebraic closure of $k((z))$. Modifying h by a suitable element in $K_{1,k((z))^{\text{alg}}}$ we may assume that it is defined over a finite extension of $k((z))$. We may replace $k((z))$ by that extension and thus assume that h is defined over $k((z))$ itself. As $K/I \cong G(k)/B(k)$ is proper, there is a $k[[z]]$ -valued point of K/I such that the induced $k((z))$ -valued point coincides with the one induced by h . Let $\tilde{h} \in G(k[[z, t]])$ be a lift of that point (which exists because $k[[z]]$ is local). Denote by $\tilde{h}_0$ and $\tilde{h}_\eta$ the images in $G(k[[t]])$ and $G(k((z))[[t]])$, respectively. As the generic points of h and $\tilde{h}$ coincide up to an element of $I_{k((z))}$, we obtain that $\tilde{h}_\eta^{-1}g_\eta\sigma(\tilde{h}_\eta) \in Iw\tau_\mu I$. Hence $\tilde{h}_0^{-1}g_0\tilde{h}_0 \in \overline{Iw\tau_\mu I}$ which proves (4). $\square$

Before proving Theorem 4 we need some preparations.

Remark 15. If $x, y, z \in \widetilde{W}$ with $x \in IyIzI$ then $x = y'z$ for some $y' \leq y$. Indeed, this follows by induction from $Is_iIzI \subseteq IzI \cup Is_izI$ for each (finite or affine) simple reflection s_i (see also [He], Lemma 4.2).

Lemma 16. *Let M be a standard Levi subgroup of G and let $x \in {}^M\widetilde{W}$. Let $y \in \widetilde{W}$. Then $y \geq wx\sigma(w)^{-1}$ for some $w \in W_M$ if and only if there are $u, v \in W_M$ with $v \leq u$ and $y \geq ux\sigma(v)^{-1}$.*

We follow the proof of [He], Proposition 3.8, a similar result for finite Weyl groups and without the σ -action (but allowing disconnected groups).

Proof. We use induction on the size of the Levi subgroup and thus may assume that the statement is true for all $M' \subsetneq M$. We use a second induction on the length $l(u)$. We write $x = ab$ with $a \in {}^M\widetilde{W} \cap \widetilde{W}^{\sigma(M)}$ and $b \in W_{\sigma(M)}$. Setting $M' = M \cap a\sigma(M)a^{-1}$ we decompose u as u_1u_2 with $u_1 \in W_{M'}$ and $u_2 \in W_{M'}$ as well as $v = v_1v_2$ with $v_i \leq u_i$ and $l(v) = l(v_1) + l(v_2)$. Note that our choice of a implies that M' is again the Levi factor of a standard parabolic subgroup. We consider two cases:

Case 1: $u_1 = v_1 = 1$. In this case u and v are in $W_{M'}$, and $x \in {}^{M'}\widetilde{W}$. If $M' \neq M$, then the assertion follows from the induction hypothesis. If $M' = M = a\sigma(M)a^{-1}$, then since $ab \in {}^M\widetilde{W}$, we have that $b = 1$. Thus $ux\sigma(v)^{-1} \geq x$ which implies the assertion.

Case 2: $u_1 \neq 1$. In this case $l(u_2) < l(u)$. By induction hypothesis, there is an $x' = u'x\sigma(u')^{-1} \leq u_2x\sigma(v_2)^{-1}$. Let $v_3 \leq v_1$ be such that $x'\sigma(v_3)^{-1}$ is the unique element of minimal length in $\{x'\sigma(v')^{-1} \mid v' \leq v_1\}$. Then $l(x'\sigma(v_3)^{-1}) = l(x') - l(v_3)$ and $x'\sigma(v_3)^{-1} \leq u_2x\sigma(v_2)^{-1}\sigma(v_3)^{-1} \leq u_2x\sigma(v)^{-1}$. We can write $x\sigma(v)^{-1} = \alpha\beta$ with $\alpha \in W_{M'}$ and $\beta \in {}^M\widetilde{W}$ (see for example Lemma 3.6 of [He]). Thus $l(u_1u_2x\sigma(v)^{-1}) = l(u_1u_2\alpha\beta) = l(u_1u_2\alpha) + l(\beta) = l(u_1) + l(u_2\alpha) + l(\beta) = l(u_1) + l(u_2\alpha\beta) = l(u_1) + l(u_2x\sigma(v)^{-1})$. As $x'\sigma(v_3)^{-1} \leq u_2x\sigma(v)^{-1}$ and $v_3 \leq v_1 \leq u_1$, we have $(v_3u')x\sigma(v_3u')^{-1} = v_3x'\sigma(v_3)^{-1} \leq ux\sigma(v)^{-1} \leq y$. $\square$

Proof of Theorem 4. We have to show that (w', μ') is the truncation of level 1 of an element of IyI for some $y \leq w\tau_\mu$ if and only if it is of the form in the theorem. The *if* part is obvious. For the other direction we use an approach which is similar to the proof of Theorem 1 to compute the truncations of level 1 occurring in the cosets IyI for y as above. We decompose y as $w_1\tau_\mu w'_1$ with $w_1, w'_1 \in W$, μ' dominant and such that the lengths of the three elements add up to that of y . Each truncation of an element of IyI already occurs in

$Iy = Iw_1\tau_{\mu'}w'_1$, and thus also in $\sigma^{-1}(w'_1)Iw_1\tau_{\mu'}$. By Remark 15, each such element is contained in $I\sigma^{-1}(\tilde{w}'_1)w_1\tau_{\mu'}I$ for some $\sigma^{-1}(\tilde{w}'_1) \leq \sigma^{-1}(w'_1)$. This is equivalent to $\tilde{w}'_1 \leq w'_1$ as I is invariant under σ . Replacing y by a smaller element we see that we may assume that $\tilde{w}'_1 = w'_1$ and that $l(\sigma^{-1}(w'_1)w_1\tau_{\mu'}) = l(w_1\tau_{\mu'}w'_1) = l(w_1) + l(\tau_{\mu'}) + l(w'_1)$. Thus we have to consider the truncation types occurring in $\sigma^{-1}(w'_1)w_1\tau_{\mu'}I$. It is enough to show that for each such type (w', μ') there is a $u \in W$ with $uw'\tau_{\mu'}u^{-1} \leq \sigma^{-1}(w'_1)w_1\tau_{\mu'}$. Indeed, then there is a $v_1 \leq \sigma^{-1}(w'_1)$ such that $v_1^{-1}uw'\tau_{\mu'}u^{-1}\sigma(v_1) \leq w_1\tau_{\mu'}w'_1$. By Lemma 16, it is furthermore enough to show the following claim.

Claim. Let (w', μ') be the truncation of level 1 of an element of $Ix\tau_{\mu'}I$ for some $x \in W$. Then there are $v \leq u \in \sigma^{-1}(W_{M_{\mu'}})$ with $uw'\tau_{\mu'}\sigma(v)^{-1} = x\tau_{\mu'}$.

We use induction to define a decreasing sequence of Levi subgroups $\tilde{M}_i \subseteq \sigma^{-1}(M_{\mu'})$, and elements $u_i \in \sigma^{-1}(W_{M_{\mu'}}) \cap W^{\tilde{M}_i}$ and $v_i \leq u_i$ with $u_ix\tau_{\mu'}\sigma(v_i)^{-1} = y_i\tau_{\mu'} \in W_{\tilde{M}_i} \cdot {}^{\mu'}\widetilde{W}$ for some y_i and such that $y_i\tau_{\mu'}I$ contains an element of truncation type (w', μ') . Recall that ${}^{\mu'}W = \sigma^{-1}(M_{\mu'}W)$. For $i = 1$ we choose $\tilde{M}_1 = \sigma^{-1}(M_{\mu'})$, and $u_1 = v_1 = 1$. For the induction step we decompose y_i as $y'_i c_i \tau_{\mu'}$ with $y'_i \in W_{\tilde{M}_i}$ and $c_i \tau_{\mu'} \in {}^{\mu'}\widetilde{W}$. Let $\tilde{M}_{i+1} = \sigma^{-1}(M_{\mu'}) \cap (c_i x_{\mu'})\sigma(\tilde{M}_i)(c_i x_{\mu'})^{-1}$. We decompose y'_i as $a_i b_i$ with $a_i \in W_{\tilde{M}_i} \cap W^{\tilde{M}_{i+1}}$ and $b_i \in W_{\tilde{M}_{i+1}}$. By σ -conjugating the element of $y_i\tau_{\mu'}I$ of truncation type (w', μ') by a_i we obtain (by Remark 15) an element of some $Ib_i c_i \tau_{\mu'}\sigma(d_i)I$ with $d_i \leq a_i$. As $u_i \in W^{\tilde{M}_i}$ and $a_i \in W_{\tilde{M}_i} \cap W^{\tilde{M}_{i+1}}$, we have that $u_{i+1} = u_i a_i \in W^{\tilde{M}_{i+1}}$ has length $l(u_i) + l(a_i)$ and hence $v_{i+1} = v_i d_i \leq u_{i+1}$. Furthermore by definition of $\tilde{M}_{i+1}$ we have that $b_i c_i \tau_{\mu'}\sigma(d_i) = y_{i+1}\tau_{\mu'} \in W_{\tilde{M}_{i+1}} \cdot {}^{\mu'}\widetilde{W}$ for some y_{i+1} . For sufficiently large n , we have that $\tilde{M}_n = \tilde{M}_{n+1}$. If we set $u = u_n y'_n$ and $v = v_n$ we have that $l(u) = l(u_n) + l(y'_n)$, hence $v \leq u$. Furthermore, by Lang's theorem and Theorem 1 (2), each element of $y_n\tau_{\mu'}I$ is K - σ -conjugate to an element of $K_1 c_n \tau_{\mu'} K_1$. Hence $c_n = w'$, which proves the claim. $\square$

The following corollary to the theorem which considers the special case $\mu = \mu'$ is analogous to results by He [He] and Wedhorn [W].

Corollary 17. $S_{w', \mu} \subseteq \overline{S_{w, \mu}}$ if and only if there is a $\tilde{w} \in \sigma^{-1}(W_{M_{\mu}})$ with $\tilde{w}^{-1}w'\tau_{\mu}\sigma(\tilde{w})x_{\mu}^{-1} \leq w$.

Proof. Recall that τ_{μ} is the unique shortest element of the affine Weyl group lying in $Kt^{\mu}K$. Especially, $y \leq w\tau_{\mu}$ with $y \in Kt^{\mu}K$ if and only if $y = w_y\tau_{\mu}$ for some $w_y \leq w$ in W . From the theorem we obtain that $S_{w', \mu} \subseteq \overline{S_{w, \mu}}$ if and only if there is a $\tilde{w} \in W$ such that $\tilde{w}^{-1}w'\tau_{\mu}\sigma(\tilde{w}) = w_y\tau_{\mu}$ for some w_y as above. Thus $\tau_{\mu}\sigma(\tilde{w}) = v\tau_{\mu}$ for some $v \in W$. As $\tau_{\mu} = x_{\mu}t^{\mu}$ we obtain $v = x_{\mu}\sigma(\tilde{w})x_{\mu}^{-1}$ and $\sigma(\tilde{w}) \in W_{M_{\mu}}$. $\square$

Recall that we have a bijection between the set $B(G)$ of σ -conjugacy classes in $G(L)$ and the set of minimal truncation types. The following proposition generalizes Corollary 3.2 of [Ha2] where the case of polarized p -divisible groups is considered. In Harashita's papers, it is used as a step in the proof of the analog of Theorem 5 whereas we first proved the theorem and now use it to deduce the proposition.

Proposition 18. Let (w, μ) and (w', μ') be the minimal truncation types associated to two σ -conjugacy classes $[b]$ and $[b']$. Then $S_{w, \mu} \subseteq \overline{S_{w', \mu'}}$ if and only if $[b] \subseteq \overline{[b']}$.

The closure of $[b']$ is a union of σ -conjugacy classes. By [RR], Theorem 3.6 a necessary condition for $[b] \subseteq \overline{[b']}$ is that $[b] \preceq [b']$, i.e. $\kappa_G(b) = \kappa_G(b')$ and $\nu_b \preceq \nu_{b'}$. It would be interesting to know if this condition is also sufficient.

Proof. Let $g \in G(k[[z]]((t)))$ such that $g_{k((z))} \in [b']$ and $g_k \in [b]$. Let $h \in G(k((z))^{\text{alg}}((t)))$ with $h^{-1}g_{k((z))}\sigma(h) \in Iw'\tau_{\mu'}I$. Here $k((z))^{\text{alg}}$ denotes an algebraic closure of $k((z))$. The

closed Schubert cell in G/I containing h is a scheme of finite type. Thus replacing h by some representative of $hI_{k((z))^{\text{alg}}}$ we may assume that h is defined over a finite extension of $k((z))$. Replacing $k[[z]]$ by its integral closure in that extension we may assume $h \in G(k((z))((t)))$. Also, as the closed Schubert cell is a proper subscheme of G/I , the class hI contains an element of $G(k[[z]]((t)))$ which we again denote by h . Then $\tilde{g} = h^{-1}g\sigma(h) \in G(k[[z]]((t)))$ with $\tilde{g}_{k((z))} \in Iw'\tau_{\mu'}I \subseteq S_{w',\mu'}$ and $\tilde{g}_k \in [b]$. Hence $\overline{S_{w',\mu'}}$ contains an element of $[b]$ and thus by Theorem 5 also the stratum $S_{w,\mu}$. $\square$

6. RELATION TO SHIMURA VARIETIES

In this section we show how Theorem 5 can be applied to the Ekedahl-Oort stratification on the moduli space $\mathcal{A}_g$ of principally polarized abelian varieties of dimension g in characteristic $p > 2$. The same considerations work for other good reductions modulo p of Shimura varieties of PEL type (except for the last corollary where one needs further input).

From an element of $\mathcal{A}_g(k)$ we obtain a polarized p -divisible group, and the considerations in Section 1.1 show that the Ekedahl-Oort stratification (as in [O1]) induced by these p -divisible groups coincides with the stratification that one obtains by considering the truncations of level 1 of the corresponding elements of $GSp_{2g}(W(k)[1/p])$. Note that by [MW] the index set for the Ekedahl-Oort stratification of $\mathcal{A}_g$ can be identified with ${}^\mu W$ where μ is the minuscule dominant element given by the Shimura datum and where W is the Weyl group of GSp_{2g} . Let S_w be the subscheme of the reduction of $\mathcal{A}_g$ corresponding to $w \in {}^\mu W$.

Remark 19. Comparisons between the Ekedahl-Oort stratification and the subdivision into Iwahori-double cosets as in Theorem 1(2) are also used in the theory of moduli spaces of abelian varieties, see for example [EG], Corollary 8.4 (iii) or [GY], 9.

Next we want to show that a p -divisible group is minimal in the sense of Oort if and only if the K - σ -conjugacy class of the element determining the Frobenius on the Dieudonné module contains a minimal element as in Section 3. We recall the notion of minimal p -divisible groups from [O3]. Let X be a p -divisible group over an algebraically closed field k and let $(\mathbf{M}, F)$ be its Dieudonné module. Let $\mathbf{N} = \mathbf{M}[1/p]$. Then X is called minimal if the endomorphisms of $(\mathbf{M}, F)$ are a maximal order in the endomorphisms of $(\mathbf{N}, F)$. There is a unique isomorphism class of minimal p -divisible groups in each isogeny class of p -divisible groups. Explicitly, if X is minimal, its Dieudonné module is isomorphic to a Dieudonné module of the following form. There is a decomposition of the rational Dieudonné module into simple summands $\mathbf{N} = \bigoplus_{i=1}^l \mathbf{N}_i$ such that $\mathbf{M} = \bigoplus_{i=1}^l \mathbf{M} \cap \mathbf{N}_i$. Let $\lambda_i = n_i/h_i$ with $(n_i, h_i) = 1$ be the slope of $\mathbf{N}_i$. Then $\mathbf{M} \cap \mathbf{N}_i$ has a basis $e_1^i, \dots, e_{h_i}^i$ such that $F(e_j^i) = e_{j+n_i}^i$. Here we use the notation $e_{j+h_i}^i = pe_j^i$.

Let now $f_j^i = e_{h_i+1-j}^i$. Let $h = \dim \mathbf{N}$. One easily checks that if we write $F = b\sigma$ for $b \in GL_h(L)$ with respect to the basis $f_1^1, \dots, f_{h_1}^1, f_1^2, \dots$, then b is contained in the Levi subgroup M given by the decomposition $\mathbf{N} = \bigoplus_{i=1}^l \mathbf{N}_i$. Furthermore, if μ denotes the M -dominant Hodge polygon of b (with respect to the choice of the upper triangular matrices as Borel subgroup), then $\mu \in \{0, 1\}^h$ is minuscule and $b = \tau_{\mu, M}$ satisfies $bI_M b^{-1} = I_M$. Hence by Lemma 9 (2) a p -divisible group is minimal if and only if it corresponds to the K - σ -conjugacy class of a minimal element.

Note that using this correspondence we see that Proposition 6.3.1 of [GHKR] (see also Remark 8) implies the main theorem of [O3] that all elements of the Ekedahl-Oort stratum of a minimal p -divisible group are isomorphic to each other.

One can use Theorem 5 to obtain

Corollary 20. *Let x be a k -valued point of S_w for some w . Let $x_0 \in \mathcal{A}_g(k)$ be a point corresponding to a minimal p -divisible group in the isogeny class of x . Then $x_0 \in \overline{S_w}$.*

This has been conjectured by Oort [O2], Conjecture 6.9 and has been shown previously by Harashita in [Ha3] using different methods. However, the strategy applied here also leads to variants without polarization, or for p -divisible groups with endomorphisms.

Proof. Let μ be the Hodge vector associated to $\mathcal{A}_g$. By the main theorem of [W] together with (6.4) of loc. cit. the closure relations between the strata S_w are known to coincide with those between the corresponding strata $S_{w,\mu}$ in the loop group of $G = GSp_{2g}$. By Theorem 5 we see that in order to prove the corollary it is enough to show that the following property holds for any pair (κ, ν) where $\kappa \in \pi_1(G)$ and $\nu \in X_*(T)_{\mathbb{Q}}/W$ are the parameters corresponding to some isogeny class of p -divisible groups: If S_w intersects the isogeny class associated to (κ, ν) then $S_{w,\mu}$ intersects the σ -conjugacy class in the loop group of G corresponding to the same pair (κ, ν) . The first condition implies that $Iw\tau_{\mu}I$ (for I defined with respect to $W(k)$) intersects the σ -conjugacy class in $G(W(k)[1/p])$ associated to (κ, ν) , the second condition is equivalent to the analogous assertion with $W(k)$ replaced by $k[[t]]$. There is a combinatorial description of these last two conditions in terms of the affine Weyl group ([GHKR], Proposition 13.3.1). In particular, the condition is independent of the choice of $W(k)$ or $k[[t]]$. $\square$

For the moduli space $\mathcal{A}_g$ of principally polarized abelian varieties of dimension g in characteristic $p > 2$ we know by [EG], Theorem 11.5 that each Ekedahl-Oort stratum which is not contained in the supersingular locus is irreducible. In particular, there is a unique generic Newton polygon in each Ekedahl-Oort stratum S_w of $\mathcal{A}_g$. Then in the same way as for the loop group we can use the above result to determine this Newton polygon.

Corollary 21. *Let ν be the generic Newton polygon in $S_w \subseteq \mathcal{A}_g$ for some $w \in {}^uW$. Then ν is the maximal element in the set of Newton polygons of minimal elements x such that $x \in \overline{S_w}$.*

Note that as long as G is split, the nonemptiness criterion [GHKR], Corollary 12.1.2 also yields a corresponding necessary condition for nonemptiness of the intersection of a given Ekedahl-Oort stratum with a given Newton stratum.

REFERENCES

- [B] R. Bédard, *On the Brauer liftings for modular representations*, J. Algebra **93** (1985), 332–353.
- [EG] T. Ekedahl, G. van der Geer, *Cycle Classes of the E-O Stratification on the Moduli of Abelian Varieties*, preprint, 2004, math.AG/0412272.
- [GHKR] U. Görtz, Th. J. Haines, R. E. Kottwitz, D. C. Reuman, *Affine Deligne-Lusztig varieties in affine flag varieties*, preprint, arXiv:0805.0045.
- [GY] U. Görtz, C.-F. Yu, *The supersingular locus in Siegel modular varieties with Iwahori level structure*, arXiv:0807.1229v1.
- [Ha1] S. Harashita, *Ekedahl-Oort strata and the first Newton slope strata*, Journal of Algebraic Geometry **16** (2007), 171–199.
- [Ha2] S. Harashita, *Configuration of the central streams in the moduli of abelian varieties*, to appear in the Asian Journal of Mathematics.
- [Ha3] S. Harashita, *Generic Newton polygons of Ekedahl-Oort strata: Oort’s conjecture*, <http://www.ms.u-tokyo.ac.jp/~harasita/gnp.ps>
- [Ha4] S. Harashita, *The supremum of Newton polygons of p -divisible groups with a given p -kernel type*, <http://www.ms.u-tokyo.ac.jp/~harasita/snp.ps>
- [HV] U. Hartl, E. Viehmann, *The Newton stratification on deformations of local G -shtuka*, preprint, <http://arxiv.org/abs/0810.0821>.
- [He] X. He, *The G -stable pieces of the wonderful compactification*, Trans. AMS **359** (2007), 3005–3024.
- [K1] R. E. Kottwitz, *Isocrystals with additional structure*, Comp. Math. **56** (1985), 201–220.
- [L] G. Lusztig, *Parabolic character sheaves. I*, Mosc. Math. J. **4** (2004), 153–179.
- [MW] B. Moonen, T. Wedhorn, *Discrete invariants of varieties in positive characteristic*, IMRN 2004, 3855–3903.
- [O1] F. Oort, *A stratification of a moduli space of abelian varieties*, in: Moduli of abelian varieties (Texel Island, 1999), 345–416, Progr. Math., 195, Birkhäuser, Basel, 2001.
- [O2] F. Oort, *Foliations in moduli spaces of abelian varieties*, J.A.M.S. **17** (2004), 267–296.
- [O3] F. Oort, *Minimal p -divisible groups*, Ann. of Math. **161** (2005), 1021–1036.

- [RR] M. Rapoport, M. Richartz, *On the classification and specialization of F -isocrystals with additional structure*, Compositio Math. **103** (1996), 153–181.
- [W] T. Wedhorn, *Specialization of F -zips*, <http://arxiv.org/abs/math/0507175>.

MATHEMATISCHES INSTITUT DER UNIVERSITÄT BONN, ENDENICHER ALLEE 60, 53115 BONN, GERMANY